

**CONTENT OF ROSMARINIC ACID IN LEAVES OF SOME SPECIES
OF LAMIACEAE AND BORAGINACEAE FAMILIES**

© **A. L. Budantsev,¹ A. L. Shavarda, N. A. Medvedeva,
N. V. Petrova, A. V. Leostrin**

Komarov Botanical Institute of RAS, St. Petersburg

¹E-mail: abudantsev@mail.ru

REFERENCES

1. Omoto T., Murakami Y., Shimomura K., Yoshihira K., Mori K., Nakashima T., Tanaka M., Ishimaru K. 1997. Caffeic acid derivatives in Lamiaceae and Boraginaceae plants. *Jap. J. Food Chem.* 4 (1): 11–16.
2. Petersen M., Abdullah Y., Benner J., Eberle D., Gehlen K., Hücherig S., Janiak V., Kim K.H., Sander M., Weitzel C., Wolters S. 2009. Evolution of rosmarinic acid biosynthesis. *Phytochem.* 70 (15–16): 1663–1679.
3. Shekarchi M., Hajimehdipoor H., Saednia S., Gohari A. R., Hamedani M. P. 2012. Comparative study of rosmarinic acid content in some plants of Labiatae family. *Pharm. Magazine.* 8 (29): 37–41.
4. Budantsev A. L., Lesiovskaya E. E. 2012. Rosmarinic acid: resources and biological activity. *Rastitelnye resursy.* 48 (3): 453–468. (In Russian)
5. Hippolyte I., Marin B., Baccou J., Jonard R. 1992. Growth and rosmarinic acid production in cell suspension cultures of *Salvia officinalis* L. *Plant Cell Rep.* 11 (3): 109–112.
6. Tada H., Murakami Y., Omoto T., Shimomura K., Ishimaru K. 1996. Rosmarinic acid and related phenolics in hairy root cultures of *Ocimum basilicum*. *Phytochem.* 42 (2): 431–434.
7. Xu H., Kim Y. K., Jin X. J., Lee S. Y., Park S. U. 2008. Rosmarinic acid biosynthesis in callus and cell cultures *Agastache rugosa* Kuntze. *J. Med. Plants.* 2 (9): 237–241.
8. Roy D., Mukhopadhyay S. 2012. Enhanced rosmarinic acid production in cultured plants of two species of *Mentha*. *Indian J. Exp. Biol.* 50 (11): 817–825.
9. Litvinenko V. I., Popova T. P., Simonjan A. V., Zoz I. G., Sokolov V. S. 1975. «Gerbstoffe» und Oxyzimtsäureabkömmlinge in Labiaten. *Planta Med.* 27 (4): 372–380.
10. Chirikova N. K., Olennikov D. N. 2008. Composition of the aerial part of *Scutellaria baicalensis*. *Chem. Nat. Compd.* 44 (3): 361–362.
11. Tóth J., Mrlianová M., Tekel'ová D., Koreňová M. 2003. Rosmarinic acid — an important phenolic active compound of lemon balm (*Melissa officinalis* L.). *Acta Fac. Pharmac. Univ. Comenianae.* 50: 139–146.
12. Chen Y., Zhu Z., Guo Q., Zhang L., Zhang X. 2012. Variation in concentrations of major bioactive compounds in *Prunella vulgaris* L. related to plant parts and phenological stages. *Biol. Res.* 45 (2): 171–175.

13. Ravn H., Pedersen M. F., Andary J., Borum C., Anthoni U., Christophersen C., Nielsen P. H. 1994. Seasonal variation and distribution of two phenolic compounds, rosmarinic acid and caffeic acid, in leaves and root-rhizomes of eelgrass (*Zostera marina* L.). *Ophelia*. 40: 51—61.