

**CONTENT OF IRIDIDS IN ELEVATED PART  
OF *TEUCRIUM POLIUM* (LAMIACEAE)**

© Yu. G. Rudakova,<sup>1</sup> O. I. Popova

Pyatigorsk Medico-Pharmaceutical Institute —  
a branch of GBOU VPO VolgGMU Russian Ministry of Health

<sup>1</sup>E-mail: Rudakovaya@mail.ru

REFERENCES

1. Rastitelnye resursy Rossii: Dikorastushchie tsvetkovyye rasteniya, ih komponentnyiy sostav i biologicheskaya aktivnost [Plant Resources of Russia: Wild flowering plants and their component composition and biological activity]. 2011. Vol. 4: Family Caprifoliaceae—Lobeliaceae. Ed. by A. L. Budantsev. St. Petersburg; Moscow. 630 p. (In Russian)
2. Rudakova Yu. G., Popova O. I. 2013. The chemical composition of grass *Teucrium polium* L. Sovremennye problemy nauki i obrazovaniya. 5: 1—8. URL: <http://www.science-education.ru/111—10725> (In Russian)
3. Rudakova Yu. G., Popova O. I. 2014. The study of phenolic compounds grass *Teucrium polium* L. Voprosy biologicheskoy, meditsinskoy i farmatsevticheskoy khimii. 3: 34—37. (In Russian)
4. Bahramikia S., Yazdanparast R. 2012. Phytochemistry and Medicinal Properties of *Teucrium polium* L. (Lamiaceae). Phytoter. Res. 26 (11): 1581—1593.
5. Dinga B., Debnath S., Banik R. 2011. Naturally occurring iridoids and secoiridoids. An updated Review. Pt 4. Chem. Pharm. Bull. 59 (7): 803—833.
6. Tundis R., Loizzo M. R., Menichini F., Statti G. A., Menichini F. 2008. Biological and pharmacological activities of iridoids: recent developments. Mini-Rev. Med. Chem. 8 (4): 399—420.
7. Rizk A. M., Hammouda F. M., Rimpler H. et al. 1986. Iridoids and flavonoids of *Teucrium polium* herb. Planta Med. 2: 87—88.
8. Fedoseeva L. V., Popov D. M. 1997. Quantitative determination of iridoids in raw motherwort. Farmatsiya. 4: 18—21. (In Russian)
9. Oganessian G. B., Galstyan A. M., Mnatsakanyan V. A. 1986. Fastigenin, harpagide, and 8-O-acetylharpagide from *Teucrium orientale*. Khimiya prirodnykh soedineniy. 22 (6): 732—733. (In Russian)
10. Popova O. I., Nikitina A. S., Makarova O. M. 2008. The study of iridoids *Dracocephalum moldavica* cultivated in conditions of Stavropol Territory. Khimiko-farmatsevticheskiy zhurnal. 42 (6): 39—42. (In Russian)