

**DYNAMICS OF DIVERSITY AND PRODUCTIVITY
OF ALAS STEPPE MEADOWS IN CENTRAL YAKUTIA**

© *M. C. Nikolaeva*,¹ *R. V. Desyatkin*

Institute for Biological Problems of Cryolithozone Siberian Branch of RAS, Yakutsk, Russia

¹E-mail: mayan34@yandex.ru

REFERENCES

1. Nikolaeva M. C., Desyatkin R. V. 2015. Dynamics of diversity and productivity of alas wet meadows in Central Yakutia. — *Rastitelnye resursy*. 51(1): 70–80. (In Russian)
2. Nikolaeva M. C., Desyatkin R. V. 2015. Dynamics of diversity and productivity of alas real meadows in Central Yakutia. — *Rastitelnye resursy*. 51(3): 328–335. (In Russian)
3. Desyatkin R. V. 1990. Specificity of soil formation in alases. — *Pochvovedenie*. 12: 5–15. (In Russian)
4. Desyatkin R. V. 2008. Pochvoobrazovaniye v termokarstovykh kotlovinakh — alasakh kriolitozony [Soil formation in thermocarst depressions — alases of cryolithozone]. Novosibirsk. 324 s. (In Russian)
5. Konspekt flory Yakutii: Sosudistyye rasyeniya [Check-list of the vascular flora of the Yakutia / Compilers L. V. Kuznetsova, V. I. Zakharova]. 2012. Novosibirsk. 282 p. (In Russian)
6. Desyatkin R. V. 2000. Grasshoppers role in biogeochemical circulation of taiga-alas landscapes. In: *Tezisy dokladov III sezda Dokuchayevskogo obshchestva pochvovedov (11–15 iyulya 2000, g. Suzdal)*. Moscow. Vol. 2. P. 74. (In Russian)