

**FORMATION OF THE SPATIAL PATTERN OF *PICEA AJANENSIS* (PINACEAE) POPULATION IN KOREAN
PINE-BROADLEAVED FOREST IN THE SOUTH OF THE RUSSIAN FAR EAST**

© *O. N. Ukhvatkina*,^{*1,2} *A. M. Omelko*,^{1,2} *A. A. Zhmerenetsky*,¹ *T. Ya. Petrenko*³

¹State Institute of Biology and Soil Sciences, Far Eastern Branch of the Russian Academy of Sciences,
Vladivostok, Russia

²Botanical Garden-Institute, Far Eastern Branch of the Russian Academy of Sciences, Vladivostok, Russia

³Far Eastern Federal University, Vladivostok, Russia

*E-mail: ukhvatkina@gmail.com

REFERENCES

1. Ricklefs R. E. 1990. Ecology. Heidelberg. 896 p.
2. Hubbel S. P. 2001. The unified neutral theory of biodiversity and biogeography. Princeton. 390 p.
3. Whittaker R. H., Levin S. A. 1977. The role of mosaic phenomena in natural communities. Theoretical population biology. 12: 117—139.
4. Hao Z., Zhang J., Song B., Ye J., Li B. 2007. Vertical structure and spatial associations of dominant tree species in an old-growth temperate forest. — For. Ecol. Manag. 252: 1—11.
5. Krestov P. V. 2003. Forest vegetation for Easternmost Russia (Russian Far East). — For. Veg. Northeast Asia. Dordrecht. P. 93—180.
6. Ukhvatkina O. N., Omelko A. M., Zhmerenetsky A. A., Gusev V. S. 2015. Natural disturbance regime as a factor of Korean pine-broadleaved forest stand shaping in the South of Russian Far East. — Zhurnal Sibirskogo federalnogo universiteta. Ser. Biologiya. 8(4): 441—458. (In Russian)
7. Omelko A. M., Ukhvatkina O. N., Zhmerenetsky A. A. 2016. Disturbance history and natural regeneration of an old-growth Korean pine-broadleaved forest in the Sikhote-Alin mountain range, Southeastern Russia. — For. Ecol. Manag. 360: 221—234.
8. Ukhvatkina O. N., Omelko A. M. 2013. Evaluation of successional processes in conifer-broadleaved forests of the Russian Far East using a population-demographic approach. — Fundamentalniye issledovaniya. 11(5): 948—953. (In Russian)
9. Liu Y. Y., Jin G. Z. 2012. Spatial distribution patterns and dynamics of four dominant tree species in a typical mixed broadleaved-Korean pine forest. In: Advances in biodiversity conservation and research in China IX. China. P. 278—296.
10. Kolesnikov B. P. 1956. Kedrovye lesa Dalnego Vostoka [Korean pine forests in the Far East]. — Trudy of DVF AN SSSR. Ser. Botanicheskaya. 2(4). Moscow; St. Petersburg. 264 p. (In Russian)

11. U s e n k o N. V. 1968. Dereviya, kustarniki i liany Dalnego Vostoka [Trees, shrubs and lianas of the Far East]. Khabarovsk. 416 p. (In Russian)
12. I s h i k a w a Y., K r e s t o v P. V., N a m i k a w a K. 1999. Disturbance history and tree establishment in old-growth *Pinus koraiensis*-hardwood forests in the Russian Far East. — J. Veg. Sci. 10: 439—448.
13. Z h a n g Y., D r o b y s h e v I., G a o L., Z h a o X., B e r g e r o n Y. 2014. Disturbance and regeneration dynamics of a mixed Korean pine dominated forest on Changbai Mountain, North-Eastern China. — Dendrochronologia. 32: 21—31.
14. L i u Y., L i F., J i n G. 2014. Spatial patterns and associations of four species in an old-growth temperate forest. — J. Plant Interact. 9(1): 745—753.
15. K o z h e v n i k o v a N. K. 2009. Dynamics of weather and climatic characteristics and ecological functions of small forest basin. — Sibiskiy ekologicheskiy zhurnal. 16(5): 693—703. (In Russian)
16. K o m a r o v a T. A., T r o f i m o v a A. D. 2010. Stages of *Pinus koraiensis* (Pinaceae) ontogenesis in Southern Sikhote-Alin mountains. — Rastitelniye resursy. 46(4): 1—15. (In Russian)
17. U k h v a t k i n a O. N., K o m a r o v a T. A., T r o f i m o v a A. D. 2010. The features of ontogenesis *Picea ajanensis* (Lindl. et Gord.) Fisch. ex Carr. in the middle part of Southern Sikhote-Alin mountains. — Vestnik Moskovskogo gosudarstvennogo universiteta lesa — Lesnoy vestnik. 3: 169—173. (In Russian)
18. S m i r n o v a O. V., B o b r o v s k i y M. V. 2001. Tree ontogeny and its reflection in the structure and dynamics of plant and soil covers. — Rus. J. Ecol. 32(3): 159—163.
19. R i p l e y B. D. 1981. Spatial Statistics. N. Y. 272 p.
20. W i e g a n d T., H e F., H u b b e l S. P. 2013. A systematic comparison of summary characteristics for quantifying point patterns in ecology. — Ecography. 36: 092—103.
21. I l l i a n J. B., P e n t t i n e n A., S t o y a n H., S t o y a n D. 2008. Statistical Analysis and Modelling of Spatial Point Patterns. Chichester. 536 p.
22. W i e g a n d T., M o l o n e y K. A. 2004. Rings, circles and null-models for point pattern analysis in ecology. — Oikos. 104: 209—229.
23. K o m a r o v a T. A. 1986. Semennoe vozobnovleniye rasteniy na svezhikh garyakh (Lesa Yuzhnogo Sikhote-Alinya) [Seed renewal of plants on the fresh burned areas (Forests of South Sikhote-Alin)]. Vladivostok. 222 p. (In Russian)
24. A l e k s e y e v V. I. 1978. Vozobnovlenie eli na vyrubkakh [Renewal of spruce on clearings]. Moscow. 130 p. (In Russian)
25. Z h a o X. H., Z h a n g C. Y., Z h e n g J. M. 2006. Correlations between canopy gaps and species diversity in broadleaved and Korean pine mixed forests. — Front. For. China. 4: 372—378.