

**SPECIES COMPOSITION AND PHYTOMASS OF THE GROUND VEGETATION IN 25-YEAR OLD
LARIX SIBIRICA (PINACEAE) ARTIFICIAL FORESTS WITH DIFFERENT STAND DENSITY**

© I. A. Goncharova

Sukachev Institute of Forest SB RAS,

Federal Research Center «Krasnoyarsk Science Center SB RAS», Krasnoyarsk, Russia

E-mail: iagoncharova007@mail.ru

REFERENCES

1. Smirnova O. V. 1994. Sinuzialnaya organizatsiya ustoichivogo lesnogo fitotsenoza [Synuzial organization of stable forest ecosystem]. In: Vostochnoyevropeiskiy shirokolistvenniy lesa. Moscow. 363 p. (In Russian)
2. Yablokov A. S. 1934. Kultura listvenitsy i ukhod za nasazhdeniyami [Artificial larch stands and caring for plantations]. Moscow. 127 p. (In Russian)
3. Timofeyev V. P. 1957. The influence of stand density on their productivity. — *Doclady TSKhA*. XXXI: 215–220. (In Russian)
4. Timofeyev V. P. 1959. Experience of larch cultivation at different planting density. — *Izvestiya TSKhA*. 2: 123–140. (In Russian)
5. Bannikova I. A. 1967. Vliyaniye drevesnoi i kustarnkovoi rastitelnosti na razvitie niznykh yarusov lesnykh biogeotsenozov [Influence of tree and shrub vegetation on the development of lower layers of forest ecosystems]. Moscow. 103 p. (In Russian)
6. Collins B. S., Pickett S. T. A. 1987. Influence of canopy opening on the environment and herb layer in a northern hardwood forest. — *Vegetatio*. 70: 3–10.
7. Økland T., Rydgen K., Økland R. H., Storaunet K. O., Rolstad J. 2003. Variation in environmental conditions among natural and managed *Picea abies* forest stands. — *Forest Ecol. Man.* 177: 17–37.
8. Cherepanova M. Iu., Ipatov V. S. 2004. The influence of canopy-depended factors on interactions of moss, grass and dwarf-shrub species in bog fir forests. — *Vestnik Sankt-Petersburgskogo universiteta*. 3(2): 10–22. (In Russian)
9. Lebedeva V. Kh., Tikhodeyeva M. Yu., Ipatov V. S. 2005. The influence of tree layer on plant soil cover species in bilberry-moss spruce forest. — *Botanicheskii Zhurnal*. 90(3): 400–410. (In Russian)
10. Lebedeva V. Kh., Tikhodeyeva M. Yu., Ipatov V. S. 2008. The effect of the tree layer on the soil cover in the bilberry aspen forest. — *Botanicheskii Zhurnal*. 93(7): 996–1010. (In Russian)

11. Zalesov S. V., Galako V. A., Predeina I. V., Pavlov A. N. 2006. Specific composition and above-ground phytomass of the live soil cover in cedar woods with different periods of age-restoration dynamics. — Izvestiya Orenburgskogo gosudarstvennogo agrarnogo universiteta. 1(9–1): 32–36. (In Russian)
12. Zasoba V. V., Danilov R. Yu. 2012. Fitotsenoticheskaya structura massivnykh lesnykh kultur Kubanskikh stepei [Phytocenotic structure of artificial forests of the Kuban steppe]. — Lesovedeniye. 1: 36–47. (In Russian)
13. Martynova M. V., Sultanova R. R. 2014. Composition and biomass of herbaceous layer in stand damaged by felling *Tilia cordata* Mill. — Agrarnyi vestnik Urala. 128(10): 59–63. (In Russian)
14. Osipov A. F., Manova S. O., Bobkova K. S. 2014. Reserves and element composition in ground cover plants in the pine forests of post-fire origin (the Komi Republic). — Rastitelnye resursy. 50(1): 3–11. (In Russian)
15. Sungurova N. R. 2015. The ground cover in cultures of pine and spruce growing in different forest-growth conditions. — Vestnik KrasGAU. 4: 153–157. (In Russian)
16. Trofimova I. A., Koshcheeva U. P., Nagimov Z. Yu., Zateeva E. A. 2015. Phytoweights live underground cover and its characteristic on a basis ecological-tsenoticheskyy scales in pine plantings of Ural educational and skilled forestry. — Agrarnyi vestnik Urala. 135(5): 55–60. (In Russian)
17. Demyanov V. A. 1977. The influence of *Larix gmelinii* (Pinaceae) on the lower layers structure in light forests at the Northern limit of its distribution (Taimyr, 72° 30'). — Botanicheskii Zhurnal. 62(9): 1333–1338. (In Russian)
18. Goncharova I. A., Sobachkin R. S. 2013. The forest stand density influence on ground layer structure in young larch artificial plantations of different density. — Vestnik KrasGAU. 3: 49–53. (In Russian)
19. Goncharova I. A., Sobachkin R. S. 2013. The ground layer structure in young artificial coniferous stands with different density. — Khvoyniye borealnoi zoni. 31(1–2): 79–83. (In Russian)
20. Goncharova I. A., Sobachkin R. S. 2015. Structure of ground vegetation in Siberian spruce cultures planted with different densities. — Russian J. Ecol. 46(4): 309–316.
21. Goncharova I. A., Sobachkin R. S. 2009. Productivity of grass-shrub layer in Scots pine plantations. — Lesovedeniye. 5: 63–68. (In Russian)
22. Ecologo-coenotic plant groups. 1990–2016. Coenotic fund of the forests of European Russia. <http://www.cepl.rssi.ru/bio/flora/ecogroup.html>
23. Shmidt V. M. 1984. Matematicheskiye metody v botanike [Mathematical methods in botany]. 288 p. (In Russian)

24. Korchagin A. A. 1976. The structure of plant communities. In: Poleyaya geobotanika. Moscow; Leningrad. Vol. 5. P. 5–320. (In Russian)
25. Melankholin P. N. 2014. Change of grass canopy under decreasing recreational load in Moscow region. — Lesovedeniye. 4: 50–55. (In Russian)
26. Moskvina I. V., Getmanets I. A. 2014. Population organization of vegetation in the subzone of southern taiga forests of the Chelyabinsk region. — Fundamentalniye issledovaniya. 12(2): 322–326. (In Russian)