

FORMATION OF SPATIAL MOSAIC OF *ABIES NEPHROLEPIS* (PINACEAE) POPULATIONS IN KOREAN PINE-BROADLEAVED FORESTS IN THE SOUTH OF RUSSIAN FAR EAST

© T. Ya. Petrenko, ^{*}1,2 A. M. Omelko, ¹A. A. Zhmerenetsky, ¹O. N. Ukhvatkina, ¹L. A. Sibirina¹

¹Federal Scientific center of the East Asia terrestrial biodiversity FEB RAS, Vladivostok, Russia

²Far Eastern Federal University, Vladivostok, Russia

*E-mail: petrenkotya@gmail.com

REFERENCES

1. Ricklefs R. E. 1990. Ecology. Heidelberg. 896 p.
2. Hubbel S. P. 2001. The unified neutral theory of biodiversity and biogeography. Princeton. 390 p.
3. Whittaker R. H., Levin S. A. 1977. The role of mosaic phenomena in natural communities. — Theoret. Pop. Biol. 12: 117–139.
4. Hao Z., Zhang J., Song B., Ye J., Li B. 2007. Vertical structure and spatial associations of dominant tree species in an old-growth temperate forest. — For. Ecol. Manag. 252: 1–11.
5. Krestov P. V. 2003. Forest vegetation for Easternmost Russia (Russian Far East). — For. Veg. Northeast Asia. Dordrecht. P. 93–180.
6. Liu Y. Y., Jin G. Z. 2012. Spatial distribution patterns and dynamics of four dominant tree species in a typical mixed broadleaved-Korean pine forest. — J. Pl. Interact. 9(1): 745–753.
7. Kolesnikov B. P. 1956. Kedrovyye lesa Dalnego Vostoka [Korean pine forest of the Far East]. Moscow. 262 p. (In Russian)
8. Usenko N. V. 1984. Derevy, kustarniki i liany Dalnego Vostoka [Trees, shrubs and lianas of the Far East]. Khabarovsk. 272 p. (In Russian)
9. Ishikawa Y., Krestov P. V., Namikawa K. 1999. Disturbance history and tree establishment in old-growth *Pinus koraiensis*-hardwood forests in the Russian Far East. — J. Veg. Sci. 10: 439–448. doi: 10.2307/3237178
10. Zhang Y., Drobyshev I., Gao L., Zhao X., Bergeron Y. 2014. Disturbance and regeneration dynamics of a mixed Korean pine dominated forest on Changbai Mountain, North-Eastern China. — Dendrochronologia. 32: 21–31.
11. Ukhvatkina O. N., Omelko A. M., Zhmerenetsky A. A., Gusev V. S. 2015. Natural Disturbances Regime as a Factor of Korean Pine-Broadleaved Forest Stand Shaping in the South of the Russian Far East. — J. Sib. Fed. Univ. Ser. biol. 4(8): 441–458. (In Russian)
12. Omelko A. M., Ukhvatkina O. N., Zhmerenetsky A. A. 2016. Disturbance history and natural regeneration of an old-growth Korean pine-broadleaved forest in the Sikhote-Alin

- mountain range, Southeastern Russia. — *For. Ecol. Manag.* 360: 221–234. doi: 10.1016/j.foreco.2015.10.036
13. Ukhvatkina O. N., Omelko A. M. 2013. Evaluation of successional processes in conifer-broadleaved forests of the Russian Far East using a population-demographic approach. — *Fundamentalniye issledovaniya*. 11(5): 948–953. (In Russian)
 14. Kozhevnikova N. K. 2009. Dynamics of weather and climatic characteristics and ecological functions of a small forest basin. — *Cont. Probl. Ecol.* 2(5): 693–703.
 15. Komarova T. A., Ukhvatkina O. N., Trophimova A. D., Kim U-San. 2010. Ontogenesis of *Abies nephrolepis* (Trautv.) Maxim. in the middle mountains of the Southern Sikhote Alin. — *Bull. Bot. Sada.-Inst. FEB RAS*. 5: 82–93. (In Russian)
 16. Smirnova O. V., Bobrovskii M. V. 2001. Tree ontogeny and its reflection in the structure and dynamics of plant and soil covers. — *Rus. J. Ecol.* 32 (3): 159–163.
 17. Illian J. B., Penttinen A., Stoyan H., Stoyan D. 2008. Statistical analysis and modelling of spatial point patterns. Chichester. 536 p.
 18. Wiegand T., He F., Hubbel S. P. 2013. A systematic comparison of summary characteristics for quantifying point patterns in ecology. — *Ecography*. 36: 092–103.
 19. Wiegand T., Moloney K. A. 2004. Rings, circles and null-models for point pattern analysis in ecology. — *Oikos*. 104: 209–229.
 20. Komarova T. A. 1986. Semennoe vozobnovleniye rasteniy na svezhikh garyakh (lesa Yuzhnogo Sikhote-Alinya) [Seed renewal of plants on the fresh burned areas (forests of the South Sikhote-Alin)]. Vladivostok. 224 p. (In Russian)
 21. Ukhvatkina O. N., Omelko A. M., Zhmerenetsky A. A., Petrenko T. Ya. 2017. Formation of the spatial pattern of *Picea ajanensis* (Pinaceae) population in Korean pine-broadleaved forest in the south of the Russian Far East. — *Rastitelnye resursy*. 53(1): 70–87. (In Russian)
 22. Ukhvatkina O. N., Omelko A. M. 2016. Life strategy of the Korean pine (*Pinus koraiensis* Sieb. et Zucc.) in the old-growth Korean pine-broadleaved forest, Southern part of the Sikhote-Alin Mountain Range. — *Vestnik Tomskogo gosudarstvennogo universiteta* [Tomsk State University Journal of Biology]. Ser. biol. 4(36):164–179. (In Russian)
 23. Grime J. P. 1977. Evidence for the existence of three primary strategies in plants and its relevance to ecological and evolutionary theory. — *Amer. Nat.* 111(982): 1169–1194.