

**AUTOMATED REMOTE EXPRESS EVALUATION OF THE DISTRIBUTION OF *PHRAGMITES AUSTRALIS* REED BEDS PROSPECTIVE FOR FUEL BIOMASS HARVESTING**

© A. A. Yanovskiy,<sup>1</sup>O. V. Sozinov<sup>\* 2,3</sup>

<sup>1</sup>Institute for Nature Management of Belarus National Academy of Sciences, Minsk, Republic of Belarus, E-mail: yanouski@ya.ru

<sup>2</sup>Komarov Botanical Institute of the RAS, St. Petersburg, Russian Federation

<sup>3</sup>Yanka Kupala Grodno State University, Grodno, Republic of Belarus

\* E-mail: ledum@list.ru

**REFERENCES**

1. Panov V. V. 2006. Vosstanovlenie torfyanykh bolot [Restoration of peat bogs]. Tomsk. 70 p. (In Russian)
2. Realizatsiya novoy kontseptsii upravleniya povtornozabolochennymi torfyanikami dlya ustoychivogo proizvodstva energii iz biomassyi (energii bolot): otchet o rabote ekspertov za 2013–2015 gg. (zaklyuch.) [Implementation of a new rewetted peatlands management concept for sustainable energy production from biomass (wetlands energy): experts report for 2013–2015 years (final)]. 2015. Institute for nature management of the NAS of Belarus; project supervisor PhD V. A. Rakovich. Minsk. 119 p. (In Russian)
3. Paludiculture – productive use of wet peatlands. Climate protection – biodiversity – regional economic benefits with contributions by 73 authors. 2016. W. Wichtmann, C. Schröder, H. Joosten. Stuttgart. 288 p.
4. Kukushkina E. E., Panov V. V. 2015. Prospects for receiving biomass of the reed (*Phragmites australis*) at natural overgrowing of the cutaway peat fields in the conditions of the Tver region. — Trudy Instorfa. 11(64): 19–31. (In Russian)
5. Rakovich V. A., Sozinov O. V., Leshchinskaya N. V., Romanovskiy Ch. A., Vikhtman V. 2016. The possibility of using plant biomass of cutover peatland «Dokudovskoe» for energy production. In: Aktualnye problemy ekologii: sb. nauch. st. po materialam XI Mezhdunar. nauch.-prakt. konf. Grodno. P. 245–247. (In Russian)
6. Strategiya sohraneniya i ratsionalnogo (ustoychivogo) ispolzovaniya torfyanikov: postanovlenie Soveta Ministrov Resp. Belarus, 30 dekabrya 2015. [The strategy of conservation and rational (sustainable) use of peatlands: Resolution of the Council of Ministers of the Republic of Belarus, 30 December 2015]. N 1111. Nac. reestr pravovykh aktov Resp. Belarus. 7 January 2016. N 5/41510. (In Russian)
7. Michael Succow Foundation. 2009. <http://www.succow-stiftung.de>

8. Ailian C., Yunbo W., Jie Z., Yanhua W. 2008. Retrieval of reed biomass based on multi-time remote sensing data: a case study on ShuangTai Estuary Nature reserve, Panjin. Remote Sensing of the Environment: 16th National Symposium on Remote Sensing of China. Proceedings of SPIE. 7123. P. 71230X.
9. Niu T., Li X., Lin H. J., Zhao Z., Dong D. R. 2011. The construction of the remote sensing estimation models for reed biomass in the lower reaches of the Tarim River. — Remote Sens. Land Res. 4: 42–45.
10. Lu S., Shimizu Y., Ishii J., Funakoshi S., Washitani I., Omasa K. 2009. Estimation of abundance and distribution of two moist tall grasses in the Watarase wetland, Japan, using hyperspectral imagery. — ISPRS J. Photogram. Remote Sens. 64(6): 674–682.
11. Luo S., Wang C., Xi X., Pan F., Qian M., Peng D., Nie S., Qin H., Lin Y. 2017. Retrieving aboveground biomass of wetland *Phragmites australis* (common reed) using a combination of airborne discrete-return LiDAR and hyperspectral data. — Int. J. App. Earth Observ. Geoinform. 58: 107–117.
12. Tuominen J., Lipping T. 2016. Spectral characteristics of common reed beds: Studies on spatial and temporal variability. — Remote Sens. 8(3): 181–199.
13. Ponyatovskaya V. M. 1964. The account of abundance and peculiarity of species arrangement in the natural plant communities. In: Polevaya geobotanika [Field geobotany]. Moscow; Leningrad. Vol. 3. P. 209–299. (In Russian)
14. Tichy L. 2002. JUICE, software for vegetation classification. — J. Veg. Sci. 13(3): 451–453.
15. Ellenberg H. 1992. Zeigerwerte der Gefäßpflanzen Mitteleuropas. Gottingen. 282 S.
16. Sozinov O. V., Rakovich V. A., Leshchinskaya N. V. 2014. Eco-coenotic characteristic of peatland «Dokudovskoe» phytocenoses on watering gradient. In: Aktualnyye problemy ekologii: materialy X Mezhdunar. nauch.-prakt. konf. In 2 parts. Part 1. Grodno. P. 46–48. (In Russian)
17. Chance K., Kurucz R. 2010. An improved high-resolution solar reference spectrum for Earth's atmosphere measurements in the ultraviolet, visible, and near infrared. — J. Quant. Spectrosc. Radiat. Transf. 111(9): 1289–1295.
18. R Core Team. 2016. R: A language and environment for statistical computing. <http://www.R-project.org/>
19. Akima H. 1991. A method of univariate interpolation that has the accuracy of a third-degree polynomial. — ACM Transactions on Mathematical Software (TOMS). 17(3): 341–366. doi: 10.1145/114697.116810
20. Akima H., Gebhardt A., Petzold T., Maechler M. 2013. akima: Interpolation of irregularly spaced data. R package version 0.5-10. <http://CRAN.R-project.org/package=akima>
21. Yanovskiy A. A. 2015. Avtomatezirovannaya otsenka sostoyaniya poverkhnosti povtorno zabolochennykh i osushennykh torfyanykh mestorozhdeniy: svidetelstvo o registratsii komputernoy

programmy v Natsionalnom tsentre intellektualnoy sobstvennosti Respubliki Belarus N 821 [Automated assessment of surface condition of rewetted and drained peatlands: a certificate of registration of the computer program in the National Center of Intellectual Property of the Republic of Belarus N 821] Received. 21.08.2015; Publ. date. 16.10.2015. (In Russian)

22. Avtomatizirovannoye tematicheskoye kartografirovaniye poverkhnosti narushennykh torfyanykh mestorozhdeniy: svidetelstvo o registratsii komputernoy programmy v Natsionalnom tsentre intellektualnoy sobstvennosti Respubliki Belarus N 816 [Automated thematic mapping of the surface of disturbed peatlands: a certificate of registration of the computer program in the National Center of Intellectual Property of the Republic of Belarus N 816]. Received. 21.08.2015; Publ. date. 16.10.2015. (In Russian)

23. Chavez P. S. 1988. An improved dark-object subtraction technique for atmospheric scattering correction of multispectral data. — *Remote Sens. Environ.* 24(3): 459–479.

24. Chavez P. S. 1989. Radiometric calibration of landsat thematic mapper multispectral images photogram. — *Eng. Remote Sens.* 55(9): 1285–1294.

25. ASTER User's Guide Part II. Level 1 Data Products (Ver. 5.1). March, 2007. Earth Remote Sensing Data Analysis Center. 66 p.

26. Iwasaki A., Fujisada H. 2005. ASTER geometric performance. — *IEEE Trans. Geosci. Remote Sens.* 43(12): 2700–2706.

27. Brockmann Consult and contributors. 2014. VISAT Version 5.0. <http://www.brockmann-consult.de/cms/web/beam/>

28. The Wetland Energy Project. 2015. Implementation of new concepts for wet peatland management for the sustainable production of biomass-based energy (wetland-energy). Newsletter N 8. December. Project registered August the 9th, 2012 N 02/12/000549. 11 p.

29. Tucker C. J. 1979. Red and photographic infrared linear combinations for monitoring vegetation. — *Remote Sens. Environ.* 8(2): 127–150. doi: 10.1016/0034-4257(79)90013-0

30. McFeeters S. 1996. The use of the Normalized Difference Water Index (NDWI) in the delineation of open water features. — *Int. J. Remote Sens.* 17: 1425–1432.

31. Courel M.-F., Chamard Ph. C., Guenegou M. C., Lerhun J., Levasseur J., Togola M. 1991. Utilisation des bandes spectrales du vert et du rouge pour une meilleure évaluation des formations végétales actives. *Téledétection et Cartographie*. Éd. AUPELF-UREF. P. 203–209.

32. seaborn: v0.7.0. January 2016. Zenodo. <http://doi.org/10.5281/zenodo.45133>

33. Conover W. J. 1999. *Practical Nonparameteric Statistics*. 3rd edition. New York. 584 p.

34. Conover W. J., Iman R. L. 1979. On multiple-comparisons procedures. Technical Report LA-7677-MS, Los Alamos Scientific Laboratory. Los Alamos, NM. 14 p.

35. Holm S. 1979. A simple sequentially rejective multiple test procedure. — *Scand. J. Statistics*. 6(2): 65–70.
36. Pohlert T. 2014. The Pairwise Multiple Comparison of Mean Ranks Package (PMCMR). R package. <http://CRAN.R-project.org/package=PMCMR>
37. Abrams M. D., Kubiske M. E. 1990. Leaf structural characteristics of 31 hardwood and conifer tree species in central Wisconsin – influence of light regime and shade-tolerance rank. — *Forest Ecol. Manag.* 31: 245–253.
38. Ustin S. L., Gamon J. A. 2010. Remote sensing of plant functional types. — *New Phytol.* 186: 795–816.
39. Hutchison K. D., Mahoney R. L., Vermote E. F., Kopp T. J., Jackson J. M., Sei A., Lisager B. D. 2009. A geometry-based approach to identifying cloud shadows in the VIIRS cloud mask algorithm for NPOESS. — *J. Atmosph. Ocean. Technol.* 26(7): 1388–1397.
40. Masek J. G., Huang C., Wolfe R., Cohen W., Hall F., Kutler J., Nelson P. 2008. North American forest disturbance mapped from a decadal Landsat record. — *Remote Sens. Environ.* 112(6): 2914–2926.
41. Zhu Z., Woodcock C. E., Olofsson P. 2012. Continuous monitoring of forest disturbance using all available Landsat imagery. — *Remote Sens. Environ.* 122(1): 75–91.
42. Vermote E., Saleous N. January. 2007. LEDAPS surface reflectance product description. Version 2.0. 21 p.
43. QGIS Development Team. 2016. QGIS Geographic Information System. Open Source Geospatial Foundation Project. <http://www.qgis.org>
44. Wichtmann W., Oehmke C., Bärtsch S., Deschan F., Malashevich U., Tanneberger F. 2013. Combustibility of biomass from wet fens in Belarus and its potential as a substitute for peat in fuel briquettes. — *Mires and Peat*. 13(6): 1–10.
45. Shaver G. R., Street L. E., Rastetter E. B., Van Wijk M. T., Williams M. 2007. Functional convergence in regulation of net CO<sub>2</sub> flux in heterogeneous tundra landscapes in Alaska and Sweden. — *J. Ecol.* 95(4): 802–817.